



National Library of Medicine

PubMed

	UI, MEU	içille	•			
PubMed	Nucleotide	Protein	Genome	Structure	PopSet	
Search PubMed	for 99246063			Go	Clear	
	Limits	Preview/Ind	lex History	Clipboard		
<u> </u>	Display Sum	many v	Save Text On	der Details	Add to Clipboard	
Entrez PubMed						
			M, Kikuno R, Hirosa		es, Protein, Nucleot	
	<u>м, мцуајиі</u> <u>О.</u>	a IV, Tanaka A, Ko	tani H, Nomura N, Oh	<u>ara</u>		
PubMed Services	complete proteins in DNA Res. 1	Prediction of the coding sequences of unidentified human genes. XIII. The complete sequences of 100 new cDNA clones from brain which code for larg proteins in vitro. DNA Res. 1999 Feb 26;6(1):63-70.				
	PMID: 102	31032; UI: 992460	63			

Related Resources

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer





National Library PubMed of Medicine

PubMed	Nucleotide	Protein	Genome	Structure	PopSet
Search PubMed	▼ for			Go	Clear
·	Limits	Preview/In	idex History	Clipboard	
	7				
· · · · · · · · · · · · · · · · · · ·	Display A	ostract:	Save Text C	order Add to Cl	pboard

Entrez PubMed

1: DNA Res 1999 Feb 26;6(1):63-70

Related Articles, Books, Protein, Nucleotide, LinkOut

PubMed Services

Prediction of the coding sequences of unidentified human genes. XIII. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro.

Nagase T, Ishikawa K, Suyama M, Kikuno R, Hirosawa M, Miyajima N, Tanaka A, Kotani H, Nomura N, Ohara O

Related Resources

Kazusa DNA Research Institute, Kisarazu, Chiba, Japan.

As a part of our cDNA project for deducing the coding sequence of unidentified human genes, we newly determined the sequences of 100 cDNA clones from a set of size-fractionated human brain cDNA libraries, and predicted the coding sequences of the corresponding genes, named KIAA0919 to KIAA1018. The sequencing of these clones revealed that the average sizes of the inserts and corresponding open reading frames were 4.9 kb and 2.6 kb (882 amino acid residues), respectively. A computer search of the sequences against the public databases indicated that predicted coding sequences of 87 genes contained sequences similar to known genes, 53% of which (46 genes) were categorized as proteins relating to cell signaling/communication, cell structure/motility and nucleic acid management. The chromosomal locations of the genes were determined by using human-rodent hybrid panels unless their mapping data were already available in the public databases. The expression profiles of all the genes among 10 human tissues, 8 brain regions (amygdala, corpus callosum, cerebellum, caudate nucleus, hippocampus, substania nigra, subthalamic nucleus, and thalamus), spinal cord, fetal brain and fetal liver were also examined by reverse transcription-coupled polymerase chain reaction, products of which were quantified by enzyme-linked immunosorbent assay.

PMID: 10231032, UI: 99246063

Dioplay Abdent T	Save Text Order L Add to Clinhoard
Lispiay Absulti	Care Text Cide: Nucleo Cipocara
•	